



Dr Brown's

natural flow®



Nipples/Teats

Read and save these instructions for future use.

BEFORE FIRST USE

Use clean hands and clean surfaces when handling. Clean the nipple/teat and then place in boiling water for 5 minutes. This is to ensure hygiene. Wash prior to use. Be sure the nipple/teat hole is clean.

BEFORE EACH USE

Pull on the nipple/teat in all directions before each use to make sure it does not tear. Inspect carefully, especially when the child develops teeth, and replace the nipple/teat immediately if it shows any signs of wear or damage, such as cracking or change of shape. Small pieces can accidentally be inhaled and cause serious injury. Clean before each use. The nipple/teat must be clean and free of obstructions.

CLEANING

Nipples/teats are dishwasher safe (top rack only). Safe for use in electric sterilizer, microwave sterilizer or boiling water.

For your child's safety and health

WARNING!

- Continuous and prolonged sucking of fluids will cause tooth decay.
- Always check food temperature before feeding.
- Throw away at the first signs of damage or weakness.
- Keep components not in use out of the reach of children.
- Never attach to cords, ribbons, laces or loose parts of clothing. The child may be strangled.
- Never use feeding nipples/teats as a pacifier/soother.
- Remove from your baby's mouth when the baby is not feeding.
- Always use this product with adult supervision.
- Glass containers may break.
- Always check the nipples/teats for signs of wear and replace when damaged.
- Do not use a microwave oven to heat the bottle as microwave ovens can create hotspots in milk/formula and/or bottle components.
- The bottle components are not toys and should not be given to children to play with or suck on.
- Do not leave the product in direct sunlight, heat or in disinfectant (sterilizing solution) for longer than recommended as this may weaken the nipple/teat.
- Change nipples/teats after approximately six weeks of use.
- When used and sterilized intensively, nipples/teats may deteriorate after six weeks.
- Do not allow a child to walk or run while using any bottle.

